

470

AN EFFECT OF 2,4-D HERBICIDE ON THE
CELL SURFACE MEMBRANE OF ONION STEM
EXPLANTS

JULY 1971 — NUMBER 13



BY

W. J. VANDERWOUDE


C. A. LEMBI

D. J. MORRE

JHRP

JOINT HIGHWAY RESEARCH PROJECT

PURDUE UNIVERSITY AND
INDIANA STATE HIGHWAY COMMISSION



Digitized by the Internet Archive
in 2011 with funding from
LYRASIS members and Sloan Foundation; Indiana Department of Transportation

Technical Paper

AN EFFECT OF 2,4-D HERBICIDE ON THE CELL SURFACE
MEMBRANE OF ONION STEM EXPLANTS

TO: J. F. McLaughlin, Director
Joint Highway Research Project

July 7, 1971

File: 9-5-3

FROM: H. L. Michael, Associate Director
Joint Highway Research Project

Project: C-36-48C

The attached Technical Paper is submitted as a report from the HPR-1 (8) research project "Research in Roadside Development", Part III, Chemical Weed Control. The paper "An Effect of 2,4-D Herbicide on the Cell Surface Membrane of Onion Stem Explants" has been authored by W. J. VanDerWoude, C. A. Lembi and D. J. Morre and is proposed for publication.

The authors plan to submit the paper to Biochemical and Biophysical Research Communications for publication. The paper is concerned with that phase of the research directed at a better understanding of the "herbicide receptor" molecules within plants. The objective of such understanding is improvement in the safe use of herbicides.

The paper is presented to the Board for approval of such publication.

Respectfully submitted,



Harold L. Michael
Associate Director

HLM:ms

cc: F. L. Ashbaucher
W. L. Dolch
W. H. Goetz
W. L. Grecco
M. J. Gutzwiller
G. K. Hallock

M. E. Harr
R. H. Harrell
M. L. Hayes
R. D. Miles
J. W. Miller
C. F. Scholer

M. B. Scott
W. T. Spencer
J. A. Spooner
N. W. Steinkamp
H. R. J. Walsh
K. B. Woods
E. J. Yoder

Technical paper

AN EFFECT OF 2,4-D HERBICIDE ON THE CELL SURFACE
MEMBRANE OF ONION STEM EXPLANTS

by

W. J. VanDerWoude

C. A. Lembi

and

D. J. Morre

Department of Botany and Plant Pathology
Purdue University

Joint Highway Research Project

File: 9-5-3

Project: C-36-48C

Prepared as Part of an Investigation

Conducted by

Joint Highway Research Project

Engineering Experiment Station

Purdue University

in cooperation with the

Indiana State Highway Commission

and the

U.S. Department of Transportation

Federal Highway Administration

The opinions, findings and conclusions expressed in this publication are those of the authors and not necessarily those of the Federal Highway Administration.

Purdue University
Lafayette, Indiana
July 7, 1971

Summary

The activity of a plasma membrane-associated glucan synthetase of onion stem was increased by addition of the auxin herbicide 2,4-D. The effect was expressed either by pre-incubating the tissue with 2,4-D or by adding the 2,4-D to the synthetase assay. 2,4-D in the synthetase assay had little or no effect with fractions from tissue grown in the presence of 2,4-D. In the in vitro assay, both the synthetase and the 2,4-D required magnesium ions for activity. The 2,4-D effect was most pronounced on the incorporation of radioactivity into polysaccharides soluble in hot water; these polysaccharides representing about 20% of the total polysaccharide synthesized in vitro.

Lembi et al. (1971) found that plasma membrane-rich fractions isolated from maize coleoptiles by low shear homogenization and differential and sucrose gradient centrifugation, specifically bind N-1-naphthylphthalamic acid (NPA), a weak auxin and inhibitor of auxin transport. In this paper, we report a response of a plasma membrane-associated enzyme of plants, glucan synthetase, to the auxin herbicide 2,4-D. The results provide further indications of specific interactions between auxins and constituents of the plant cell surface.

Methods and Materials

Plasma membrane fractions were isolated from onion (Allium sepa) stem according to the procedure of Lembi et al. (1971). Approximately 15 g of onion stem explants (Morré, 1970) were homogenized in 35 ml of a freshly prepared medium consisting of 0.5 M sucrose, 0.1 M K_2HPO_4 , 1 mM dithiothreitol and 0.02 M EDTA in coconut milk as the solvent and adjusted to pH 6.5 with NaOH. Nuclei and other endogenous cellular components of the coconut milk were removed by high speed centrifugation (100,000 g, 90 min) before preparing the medium. Homogenates were prepared in the cold using a mechanized razor blade chopper (Morré, 1971). Cell walls and unbroken cells were removed by filtration through a single layer of miracloth (Chicopee Mills, New York), and the filtered homogenate was centrifuged for 30 min at 10,000 g (Spinco L2-65B, SW 50.1 rotor) to remove nuclei, plastids, mitochondria and large membrane fragments. The supernatant containing microsomes and the majority of the cytoplasmic membrane fragments was then sedimented at 44,000 g onto a two-layered cushion consisting of

1. The first part of the report is a general introduction to the subject.

2. The second part is a detailed description of the methods used in the study.

3. The third part is a discussion of the results of the study.

4. The fourth part is a conclusion and a list of references.

5. The fifth part is a list of figures and tables.

6. The sixth part is a list of appendices.

7. The seventh part is a list of footnotes.

8. The eighth part is a list of acknowledgments.

1. Introduction

The purpose of this study is to investigate the effects of the treatment.

The study was conducted in a laboratory setting.

The results of the study are presented in the following sections.

The first section describes the methods used in the study.

The second section describes the results of the study.

The third section discusses the implications of the study.

The fourth section concludes the study.

The fifth section lists the references.

The sixth section lists the figures and tables.

The seventh section lists the appendices.

The eighth section lists the footnotes.

The ninth section lists the acknowledgments.

The tenth section lists the references.

The eleventh section lists the figures and tables.

The twelfth section lists the appendices.

1.5 M and 2.0 M sucrose in coconut milk to which 1 mM dithiothreitol had been added. The supernatant was removed and replaced by a step gradient of 0.5, 0.8, 1.1, 1.2 and 1.3 M sucrose in coconut milk + 1 mM dithiothreitol. The gradient was then centrifuged for 90 min at 100,000 g. Plasma membranes were recovered from the 1.1/1.2 M interface of the gradient, resuspended in homogenization medium and pelleted for 20-30 min at 100,000 g. The plasma membrane fragments were identified by electron microscopy after differential staining with a mixture of phosphotungstic acid and chromic acid which specifically and characteristically stains the plant cell membrane (Morré et al., 1970; Lembi et al., 1971, in preparation).

The glucan synthetase assay is adapted from that of Ordin and Hall (1967) and is based on the incorporation of isotopically labeled sugars from appropriate sugar nucleotides into polysaccharides of varying solubility. The substrate concentration was increased to saturate the substrate requirements for the synthetases involved with glucose incorporation into lipid-soluble compounds and intermediates (Pinsky and Ordin, 1969). This activity of onion stem is characterized by a high affinity for UDP-glucose ($K_m = 2.5 \times 10^{-8}$ M) and competes with the glucan synthetases at low substrate concentrations (Lembi et al., in preparation). As shown previously by Ordin and Hall (1968) and Feingold et al. (1958) predominantly beta 1,3 linkages are produced from UDP-glucose at these high substrate concentrations.

Fractions for assay were resuspended in 0.1 M Tris-HCl containing 4 mM disodium EDTA and 1 mM dithiothreitol, final pH 8.0, at a concentration of 1-4 mg protein/ml. The assay medium contained 0.3 μ moles UDP-glucose- C^{14} (227 μ C/ μ mole), 200 μ moles unlabeled UDP-glucose, 10 μ moles Tris-HCl,

pH 8.0, 0.4 μ moles disodium EDTA, 0.1 μ moles dithiothreitol, 4 μ moles $MgCl_2$, 2 μ moles cellobiose (as the acceptor) and an aliquot of the fraction to provide 50-200 μ g protein in a final volume of 200 μ l. Incubations were for 15 min at 25°.

Prior to fractionation of the polysaccharide synthesized into hot water soluble-, lipid soluble-, sodium hydroxide soluble- and sodium hydroxide insoluble-constituents, cellulose powder (Sigmacell, Sigma Chemical Company, St. Louis, Missouri), 1-2 mg, was added to each assay tube. The mixture was then extracted 3 times, 5 min each, with 1 ml aliquots of water at 95-100°. Insoluble polysaccharides were recovered after each extraction by centrifugation at 25,000 g for 12 min. Hot water insoluble polysaccharides were recovered from the hot water extracts by addition of 10 mg ficol (mol wt. 400,000) followed by precipitation from cold 70% ethanol. The hot water insoluble residue was extracted once with 1 ml of methanol-chloroform (2:1, v/v) and once with 1 ml of absolute methanol. The methanol-chloroform and methanol extracts were combined to provide a measure of the synthesis of lipid-soluble sugar derivatives and intermediates (Pinsky and Ordin, 1969). After removal of lipids, the residue was then extracted twice with 0.25 ml aliquots of hot 1 N sodium hydroxide for 5 min followed by 0.75 ml of cold water. The combined alkali extracts (including the water wash) were evaporated to dryness for direct determination of radioactivity on planchets. The alkali insoluble residue is the "cellulose" fraction. Self absorption corrections were determined and applied to each of the fractions.

Results

The specific activity of a marker enzyme UDP-glucose: glucose glucosyltransferase (glucan synthetase), concentrated in plasma membrane fractions isolated from onion stem explants (Lembi et al., in preparation), is markedly increased by incubating the explants in the presence of the synthetic plant growth regulator 2,4-D (Table I). With tissue incubated for 12-16 hours in the presence or absence of $5\mu\text{M}$ 2,4-D, the specific activity of the glucan synthetase of plasma membranes isolated from the 2,4-D-incubated tissue was increased approximately 25% over that of plasma membranes from control tissues incubated in the absence of 2,4-D.

To demonstrate that a similar response results from the in vitro addition of the auxin, plasma membranes were isolated from tissues incubated in the presence or absence of $5\mu\text{M}$ 2,4-D. Each of the preparations was then assayed in the presence or absence of $5\mu\text{M}$ 2,4-D added to the enzyme assay medium (Table II). A consistent stimulation of 7 to 15% was observed in the synthesis of hot water insoluble glucans when 2,4-D was added to plasma membranes isolated from control tissues. With plasma membranes isolated from 2,4-D incubated tissues, incubation with 2,4-D had little or no additional effect on the glucan synthetase activity (Table II).

The products of the synthetase assay are heterogeneous and not all fractions are affected similarly by the presence of 2,4-D in the enzyme assay. After fractionation, the largest 2,4-D response of the hot water insoluble fraction is expressed in the fraction insoluble in HOT NaOH; the fraction

The first part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The second part of the paper is devoted to a discussion of the experimental results obtained in the study of the structure of the atom. It is shown that the experimental results are in good agreement with the theoretical predictions of the theory of the structure of the atom.

The third part of the paper is devoted to a discussion of the applications of the theory of the structure of the atom. It is shown that the theory of the structure of the atom has many important applications in the field of physics and chemistry. The fourth part of the paper is devoted to a discussion of the future of the theory of the structure of the atom. It is shown that the theory of the structure of the atom is still a very active field of research and that many new results are expected in the future.

soluble in hot NaOH showing much less of an effect (Table III). There was no effect of 2,4-D on glucose incorporation into the lipid soluble fraction. The glucan fraction soluble in hot water, a fraction not included in data of Tables I and II, showed the greatest stimulation (30%) by 2,4-D added to the glucan synthetase assay.

The plasma membrane-associated glucan synthetase was markedly stimulated by magnesium ions (Table IV) and the 2,4-D-induced stimulation of activity was not observed in the absence of magnesium ions. In the absence of magnesium ions, 2,4-D was without effect or inhibited the activity of the glucan synthetase. Again, the greatest stimulation was shown by the hot water soluble fraction rather than the hot water insoluble fraction (30% vs. 11% stimulation by 2,4-D).

A comparison of the relative effectiveness of in vitro and in vivo incubations with 2,4-D on the production of the hot water soluble glucan by isolated plasma membranes is given in Table V. In 3 experiments, a stimulation of 30 to 65% resulted from the inclusion of 2,4-D in the glucan synthetase assay with plasma membranes isolated from tissue incubated in the absence of 2,4-D. The inclusion of 2,4-D in the glucan synthetase assay of plasma membranes isolated from tissue incubated in the presence of 2,4-D had no additional effect except in Expt. II of Table V where incubation of the tissue with 2,4-D was not effective (-15% stimulation). In this experiment, addition of 2,4-D to the synthetase assay resulted in an apparent stimulation of 80% in the synthesis of hot water soluble glucans.

Discussion

A 50% enhancement of glucan synthetase activities by auxin pretreatment was previously reported for pea epicotyls (Abdul-Baki and Ray, 1971); an activity localized within the Golgi apparatus of this tissue by Ray et al. (1969). A similar enhancement, although not as great, appears to be associated with the plasma membrane-associated glucan synthetase of onion stem. Additionally, we show that 2,4-D added to the isolated plasma membrane fragments ~~increases~~ the specific activity of the glucan synthetase when the plasma membrane fragments are isolated from tissues which have not been pre-treated with auxin. Except for one experiment (Expt. II of Table V), 2,4-D added to plasma membranes isolated from tissues incubated with 2,4-D resulted in no additional stimulation. Although we do not rule out an effect of 2,4-D on the synthesis of glucan synthetase in vivo, at least a significant portion of the stimulation appears to result from a direct effect on the plasma membrane-associated synthetase.

The in vitro response to 2,4-D is most marked with the hot water soluble polysaccharide fraction, a fraction specifically enhanced by auxin treatment of growing tissues (Bonner, 1961; Ray, 1962; Morrè and Olson, 1965). Here, the response to 2,4-D added in vitro to the glucan synthetase assay was equivalent to results obtained with plasma membranes isolated from tissues incubated 12 to 16 hours in the presence of 2,4-D. Although related to an artificial system of wall fractionation (Albersheim, 1965), the results are important in that a response to auxin has been obtained by adding the auxin to an in vitro system.

References

1. Abdul-Baki, A. A. and Ray, P. M., *Plant Physiol.* 47, 537 (1971).
2. Albersheim, P. In Plant Biochemistry (J. Bonner and J. Varner, eds) 290-321, New York, Academic Press, 1965.
3. Bonner, J., In Plant Growth Regulation (R. H. Klein, ed) 307-320, Ames, Iowa State University Press, 1961.
4. Feingold, D. S., Neufeld, E. F. and Hassid, W. Z., *J. Biol. Chem.* 233, 703 (1958).
5. Lembi, C. A., Morré, D. J., St.-Thompson, K. and Hertel, R., *Planta* (1971).
6. Morré, D. J., *Plant Physiol.* 45, 791 (1970).
7. Morré, D. J., *Methods Enzymol.* 22, 130 (1971).
8. Morré, D. J. and Olson, A. C., *Can. J. Bot.* 43, 1003 (1965).
9. Morré, D. J., Roland, J.-C., and Lembi, C. A., *Proc. Ind. Acad. Sci.* for 1969 79, 96 (1970).
10. Ordin, L. and Hall, M. A., *Plant Physiol.* 42, 205 (1967).
11. Ordin, L. and Hall, M. A., *Plant Physiol.* 43, 473 (1968).
12. Pinsky, A. and Ordin, L., *Plant Cell Physiol.*, 10, 771 (1969).
13. Ray, P. M., *Amer. J. Bot.* 49, 920 (1962).
14. Ray, P. M., Shiningier, T. L. and Ray, M. M., *Proc. Nat. Acad. Sci. U.S.* 64, 605 (1969).

THEORY

Let $f(x)$ be a function defined on the interval $[a, b]$. The function $f(x)$ is said to be continuous at the point x_0 if for every $\epsilon > 0$ there exists a $\delta > 0$ such that for all x in the interval $[a, b]$ satisfying $|x - x_0| < \delta$, the inequality $|f(x) - f(x_0)| < \epsilon$ holds. If the function $f(x)$ is continuous at every point of the interval $[a, b]$, then it is said to be continuous on the interval $[a, b]$.

Let $f(x)$ be a function defined on the interval $[a, b]$. The function $f(x)$ is said to be differentiable at the point x_0 if the limit $\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$ exists. If the function $f(x)$ is differentiable at every point of the interval $[a, b]$, then it is said to be differentiable on the interval $[a, b]$.

Let $f(x)$ be a function defined on the interval $[a, b]$. The function $f(x)$ is said to be integrable on the interval $[a, b]$ if the limit $\lim_{n \rightarrow \infty} \sum_{k=1}^n \Delta x_k f(\xi_k)$ exists, where Δx_k is the width of the subinterval $[x_{k-1}, x_k]$ and ξ_k is a point in the subinterval $[x_{k-1}, x_k]$. If the function $f(x)$ is integrable on the interval $[a, b]$, then the integral $\int_a^b f(x) dx$ is defined as the limit $\lim_{n \rightarrow \infty} \sum_{k=1}^n \Delta x_k f(\xi_k)$.

Let $f(x)$ be a function defined on the interval $[a, b]$. The function $f(x)$ is said to be convex on the interval $[a, b]$ if for any two points x_1 and x_2 in the interval $[a, b]$ and any λ in the interval $[0, 1]$, the inequality $f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$ holds. If the function $f(x)$ is convex on the interval $[a, b]$, then it is said to be a convex function on the interval $[a, b]$.

Let $f(x)$ be a function defined on the interval $[a, b]$. The function $f(x)$ is said to be concave on the interval $[a, b]$ if for any two points x_1 and x_2 in the interval $[a, b]$ and any λ in the interval $[0, 1]$, the inequality $f(\lambda x_1 + (1 - \lambda)x_2) \geq \lambda f(x_1) + (1 - \lambda)f(x_2)$ holds. If the function $f(x)$ is concave on the interval $[a, b]$, then it is said to be a concave function on the interval $[a, b]$.

TABLE I

IN VIVO STIMULATION OF PLASMA MEMBRANE-ASSOCIATED GLUCAN SYNTHETASE BY
2,4-D PRETREATMENT OF ONION STEM TISSUE

Onion stem explants (Morre, 1970) were incubated for 12-16 hours at $25^{\circ} \pm 5^{\circ}\text{C}$ in $5\mu\text{M}$ 2,4-D. Plasma membrane fractions were isolated by the method of Lembi *et al.* (1971). Units of specific activity of the isolated fractions are m moles glucose incorporated into total hot water-insoluble polysaccharide/hr/mg protein.

Tissue preincubation	Specific activity			
	Expt. I	Expt. II	Expt. III	Expt. IV
None	319	321	334	324
$5\mu\text{M}$ 2,4-D	410	391	456	410
Δ 2,4-D %	29	22	19	26

IN VITRO 2,4-D STIMULATION OF PLASMA MEMBRANE-ASSOCIATED GLUCAN SYNTHETASE
COMPARING ONION STEM TISSUE PRETREATED $\pm$ 2,4-D

Plasma membrane fractions were isolated from stem explants and units of specific activity are as described in Table 1. 2 μ M 2,4-D was then added to the enzyme assay to test for in vitro stimulation of the membrane-associated synthetase.

Auxin added							
Tissue	Synthetase	Specific activity					
incubation	assay	Expt. II	Expt. III	Expt. IV	Expt. V	Expt. VI	Ave.
None	None	321	304	324	342	377	350
	5 μ M 2,4-D	344	443	373	366	414	380
	Δ 2,4-D, %	+7	+15	+15	+7	+10	+11
5 μ M 2,4-D	None	391	450	410	---	---	420
	5 μ M 2,4-D	401	403	393	---	---	426
	Δ 2,4-D, %	+3	+5	-4	--	--	+1

TABLE III

SUBFRACTIONATION OF POLYSACCHARIDES OF THE IN VITRO ASSAY AND EFFECT OF 2,4-D
ON INCORPORATION OF RADIOACTIVITY INTO VARIOUS SUBFRACTIONS

Onion stem explants were incubated 12-16 hours at 25° in the absence of 2,4-D. Plasma membrane fractions were isolated and glucan synthetase activity was assayed with 5 μ M 2,4-D added directly to the enzyme assay. Details of the polysaccharide fractionation are given in the text.

		mpmoles glucose incorporated/hr/mg protein			
Auxin added to synthetase assay	Expt.	Fractions soluble in			Hot NaOH
		Hot water	CHCl ₃ -MeOH	Hot NaOH	Insoluble
None	III	35	21	104	230
	V	36	22	133	214
	VI	37	14	147	230
	Ave.	70	20	130	240
5 μ M 2,4-D	III	123	19	126	317
	V	96	21	133	233
	VI	50	19	143	271
	Ave.	90	20	135	274
Δ 2,4-D, %		+30	0	+4	+14

TABLE IV

EFFECT OF MAGNESIUM IONS ON PLASMA MEMBRANE-BOUND GLUCAN SYNTHETASE AND THE
RESPONSE TO 2,4-D

Plasma membrane fractions were isolated from stem explants and units of specific activity are as described in Table III. Average of 2 experiments.

Auxin added to synthetase assay	Synthetase assay medium	mpmoles glucose incorporated/hr/mg protein	
		Hot water soluble	Hot water insoluble
None	-Mg ⁺⁺	61	80
	Complete	85	363
5 μ M 2,4-D	-Mg ⁺⁺	56	81
	Complete	110	404

TABLE V

COMPARISON OF *IN VITRO* AND *IN VIVO* STIMULATION BY 2,4-D OF THE SYNTHESIS OF HOT WATER SOLUBLE POLYSACCHARIDES BY THE PLASMA MEMBRANE-ASSOCIATED GLUCAN SYNTHETASE

Plasma membrane fractions were isolated from stem explants incubated as described in Table 1. The unit of specific activity is μ moles glucose incorporated into total hot water soluble polysaccharide/hr/mg protein.

Auxin added					
Tissue incubation	Synthetase assay	Specific activity			
		Expt. II	Expt. III	Expt. IV	Ave.
None	None	85	83	103	90
	5 μ M 2,4-D	120	111	139	134
	Δ 2,4-D, %	+45	+64	+34	+49
5 μ M 2,4-D	None	72	115	190	126
	5 μ M 2,4-D	129	112	135	142
	Δ 2,4-D, %	+30	-3	-3	+12
Δ Tissue incubation with 2,4-D, %		-15	+39	+35	+40

